I am under anticoagulation treatment with Rivaroxaban-Teva.

Name

Address

Date of birth

Weight

Other medicines/conditions

In case of emergency, please notify:

Doctor's name

Doctor's phone

Please also notify:

Name

Phone

Relationship

Document Number: Riv-SG-NP-00001 Expiration Date: March 2028

Rivaroxaban-Teva

Rivaroxaban

PATIENT ALERT CARD

- Rivaroxaban-Teva 2.5 mg ☐
- Rivaroxaban-Teva 10 mg ☐
- Rivaroxaban-Teva 15 mg ☐
- Rivaroxaban-Teva 20 mg ☐

✓ Tick the prescribed dose

- **Keep this card with you at all times**
- **Present this card to every physician or dentist prior to treatment**

teva

Information for health care providers:
INR values should not be used as they are not a dependable measure of the anticoagulant activity of Rivaroxaban-Teva.

This document has been approved by HSA as of 2 Feb 2026.

What Should I Know About Rivaroxaban-Teva ?

- Rivaroxaban-Teva thins the blood, which prevents you from getting dangerous blood clots.
- Rivaroxaban-Teva must be taken exactly as prescribed by your doctor. To ensure optimal protection from blood clots, never skip a dose.
- You must not stop taking Rivaroxaban-Teva without first talking to your doctor as your risk of blood clots may increase.
- Tell your healthcare provider about any other medicines you are currently taking, took recently or intend to start taking, before you start Rivaroxaban-Teva.
- Tell your healthcare provider that you are taking Rivaroxaban-Teva before any surgery or invasive procedure.

When Should I Seek Advice from my Healthcare Provider?

- When taking a blood thinner such as Rivaroxaban-Teva, it is important to be aware of its possible side effects.
- Bleeding is the most common side effect. Do not start taking Rivaroxaban-Teva if you know you are at risk of bleeding, without first discussing this with your doctor.
- Tell your healthcare provider straight away if you have any signs or symptoms of bleeding such as the following:
  - Pain
  - Swelling or discomfort
  - Headache, dizziness or weakness
  - Unusual bruising, nosebleeds, bleeding of gums, cuts that take a long time to stop bleeding
  - Menstrual flow or vaginal bleeding that is heavier than normal
  - Blood in your urine which may be pink or brown, red or black stools
  - Coughing up blood, or vomiting blood or material that looks like coffee grounds

How Do I Take Rivaroxaban-Teva ?

- To ensure optimal protection, Rivaroxaban-Teva
  - 2.5 mg can be taken with or without food
  - 10 mg can be taken with or without food
  - 15 mg must be taken with food
  - 20 mg must be taken with food

关于 Rivaroxaban-Teva 我应该知道些什么?

- Rivaroxaban-Teva 稀释血液以防止发生危险性的血栓
- Rivaroxaban-Teva 必须严格按照医生的处方服用。为了最大程度预防血栓，切勿漏服剂量
- 未事先经医生同意，切勿停止服用 Rivaroxaban-Teva，因为形成血栓的风险可能会增加
- 在开始服用 Rivaroxaban-Teva 之前，请告知医护人员您正在服用、最近服用或打算开始服用的任何其他药物
- 在进行任何手术或侵入性操作之前，请告知医护人员您正在服用 Rivaroxaban-Teva.

我应该在什么时候向医护人员寻求建议?

- 服用血液稀释剂如 Rivaroxaban-Teva 时，了解其可能的药物副作用 非常重要
- 出血是最常见的副作用。如果您知道自己有出血风险，在没有事先与医生讨论的情况下，切勿开始服用 Rivaroxaban-Teva
- 如果您出现任何如下出血体征或症状，请立即告知医护人 员:
  - 疼痛
  - 肿胀或不适
  - 头痛, 晕眩或虚弱
  - 不寻常的瘀青、鼻出血、牙龈出血、伤口止血时间长
  - 流量比往常多的月经或阴道出血
  - 粉红色或褐色的尿液，红色或黑色的粪便
  - 咳嗽带血，呕血或呕出类似咖啡渣的物质

我应该怎么服用Rivaroxaban-Teva?

- 为了达到最理想的保护效果，Rivaroxaban-Teva
  - 2.5 mg 的剂量可在用餐前或用餐后服用
  - 10 mg 的剂量可在用餐前或用餐后服用
  - 15 mg 的剂量必须在用餐后服用
  - 20 mg 的剂量必须在用餐后服用

Maklumat Yang Perlu Saya Tahu Tentang Rivaroxaban-Teva?

- Rivaroxaban-Teva ialah ubat cair darah yang boleh mencegah darah beku yang berbahaya.
- Rivaroxaban-Teva mesti diambil seperti yang diarahkan oleh doktor anda. Untuk memastikan perlindungan optimum daripada darah beku, jangan tinggalkan dos yang perlu diambil.
- Jangan hentikan pengambilan Rivaroxaban-Teva tanpa berbincang dengan doktor anda terlebih dahulu kerana risiko darah beku berlaku boleh meningkat.
- Beritahu penyedia penjagaan kesihatan anda tentang sebarang ubat yang sedang anda ambil, ubat yang diambil baru-baru ini atau ubat yang anda berhasrat untuk mula ambil, sebelum anda mulakan pengambilan Rivaroxaban-Teva
- Beritahu penyedia penjagaan kesihatan anda bahawa anda mengambil Rivaroxaban-Teva sebelum pembedahan atau prosedur invasif.

Bilakah Saya Harus Dapatkan Nasihat Daripada Penyedia Penjagaan Kesihatan Saya?

- Apabila mengambil ubat cair darah seperti Rivaroxaban-Teva, kesedaran tentang kesan sampingan ubat tersebut adalah amat penting
- Pendarahan ialah kesan sampingan yang paling biasa. Jangan mula mengambil Rivaroxaban-Teva jika anda tahu anda berisiko mengalami pendarahan tanpa berbincang dengan doktor terlebih dahulu.
- Beritahu penyedia penjagaan kesihatan anda dengan segera jika anda mengalami tanda atau gejala pendarahan seperti berikut:
  - Sakit
  - Bengkak atau rasa tidak selesa
  - Sakit kepala, pening atau rasa lemah
  - Lebam, hidung atau gusi berdarah yang tidak normal, pendarahan daripada luka yang mengambil masa yang lama untuk berhenti.
  - Haid atau pendarahan faraj yang lebih banyak daripada biasa.
  - Air kencing berdarah yang boleh berwarna merah jambu atau coklat.
  - Najis berwarna merah atau hitam.
  - Batuk darah, muntah darah atau muntah serupa hampas kopi.

Bagaimanakah Saya Harus Mengambil Rivaroxaban-Teva?

- Untuk memastikan perlindungan optimum, Rivaroxaban-Teva
  - 2.5 mg boleh diambil dengan atau tanpa makanan
  - 10 mg boleh diambil dengan atau tanpa makanan
  - 15 mg mesti diambil dengan makanan
  - 20 mg mesti diambil dengan makanan

Document Number: Riv-SG-NP-00001 Expiration Date: March 2028